

Summary of 4 March Organization Meeting

Participants

Mark Cohen, NOAA ARL
David Gay, NADP MDN
Charley Driscoll, Syracuse University
David Krabbenhoft, USGS
David Evers, BRI
Richard Haeuber, EPA
David Schmeltz, EPA
Colleen Haney, EPA
Tim Sharac, EPA
Christine Negra, Consultant

By phone:

Richard Artz, *NOAA ARL*Gregory Masson, *USFWS*Kathy Fallon Lambert, *Ecologic*

Purpose

- Review information resources
- Identify potential intensive and approaches for cluster sites

Activities


- Identified and characterized existing monitoring sites
- Identified potential intensive sites
- Discussed approach to cluster sites

Site Selection Criteria

- Distribution
- 2. Sensitivity to mercury deposition
- Proximity to sources (elevated loading)
- 4. Previous studies and monitoring
- 5. Conservation concerns
- 6. Region around sites

Prior to meeting, four separate individuals/groups selected 20 potential intensive monitoring sites

- Sites discussedCountry divided into subregions
 - Southeast
 - Northeast
 - Mid-Atlantic
 - Ohio River Valley
 - Mid-West
 - West



Considerations

- High deposition/deposition changeOhio Valley
- Sensitive regions Southeast, Northeast, Upper Midwest
- Comprehensive monitoring
 - Everglades
 - Chesapeake Bay
 - ELA
 - Devil's Lake
 - Huntington Forest

Issues

- What is a site? e.g., Chesapeake Bay
- What are the regions?
- Important area, but limited intensive sites
- Important site for testing atmosphere and ecosystem models
- Scope of mercury contamination (e.g., mining, reservoirs)
- Reference site (?)
- Number of compelling intensive sites?
- Available cluster sites